Index of Authors

ALT, C. *see* KAGE, H.

ANWAR, M. R., MCKENZIE, B. A. and HILL, G. D. Phenology and growth response to irrigation and sowing date of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate, 273

ANWAR, M. R., MCKENZIE, B. A. and HILL, G. D. The effect of irrigation and sowing date on crop yield and yield components of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate, 259

ANWAR, M. R., MCKENZIE, B. A. and HILL, G. D. Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate, 285

APPENDINO, M. A. and SLAFER, G. A. Earliness per se and its dependence upon temperature in diploid wheat lines differing in the major gene *Eps-A^m 1* alleles, 149

AYAD, G. *see* BARI, A.

BADU-APRAKU, B. *see* KAMARA, A. Y.

BANTERNG, P., PATANOTHAI, A., PANNANGPETCH, K., JOGLOY, S. and HOOGENBOOM, G. Seasonal variation in the dynamic growth and development traits of peanut lines, 51

BARI, A., MARTIN, A., BOULOUHA, M., GONZALEZ-ANDUJAR, J. L., BARRANCO, D., AYAD, G. and PADULOSI, S. Use of fractals and moments to describe olive cultivars, 63

BARRANCO, D. *see* BARI, A.

BERGER, P. H. *see* HAINES, M. M.

BERRY, P. M., SPINK, J. H., GAY, A. P. and CRAIGON, J. A comparison of root and stem lodging risks among winter wheat cultivars, 191

BOULOUHA, M. *see* BARI, A.

BRENES, A. *see* CENTENO, C.

BROWNE, R. A., WHITE, E. M. and BURKE, J. I. Effect of nitrogen, seed rate and plant growth regulator (chlormequat chloride) on the grain quality of oats (*Avena sativa*), 249

BURKE, J. I. *see* BROWNE, R. A.

BUTTERY, P. J. *see* KIM, E. J.

CENTENO, C., VIVEROS, A., BRENES, A., LOZANO, A. and DE LA CUADRA, C. Effect of several germination conditions on total P, phytate P, phytase, acid phosphatase activities and inositol phosphate esters in spring and winter wheat, 313

CHEN, C. L. *see* JENG, T. L.

CIAVARELLA, T. A. *see* MCFARLANE, N. M.

CRAIGON, J. *see* BERRY, P. M.

DE LA CUADRA, C. *see* CENTENO, C.

DHANOA, M. S. *see* KIM, E. J.

DWIVEDI, B. S. *see* YADAV, R. L.

EIMERT, K., REUTTER, G. and STROLKA, B. Fast and reliable detection of doubled-haploids in *Asparagus officinalis* by stringent RAPD-PCR, 73

FELLMAN, J. K. *see* HAINES, M. M.

FREEMAN, S. E. *see* LUTMAN, P. J. W.

GAY, A. P. *see* BERRY, P. M.

GONZALEZ-ANDUJAR, J. L. *see* BARI, A.

HAINES, M. M., SHIEL, P. J., FELLMAN, J. K. and BERGER, P. H. Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETR1, 333

HAKALA, K. and PAHKALA, K. Comparison of central and northern European winter rye cultivars grown at high latitudes, 169

HILL, G. D. *see* ANWAR, M. R.

HOOGENBOOM, G. *see* BANTERNG, P.

IBIKUNLE, O. *see* KAMARA, A. Y.

JENG, T. L., WANG, C. S., CHEN, C. L. and SUNG, J. M. Effects of grain position on the panicle on starch biosynthetic enzyme activity in developing grains of rice cultivar Tainung 67 and its NaN_3-induced mutant, 303

JENNER, C. F. *see* ZAHEDI, M.

JOGLOY, S. *see* BANTERNG, P.

KAGE, H., ALT, C. and STÜTZEL, H. Aspects of nitrogen use efficiency of cauliflower I. A simulation modelling based analysis of nitrogen availability under field conditions, 1

KAGE, H., ALT, C. and STÜTZEL, H. Aspects of nitrogen use efficiency of cauliflower II. Productivity and nitrogen partitioning as influenced by N supply, 17

KAMARA, A. Y., KLING, J. G., MENKIR, A. and IBIKUNLE, O. Agronomic performance of maize (*Zea mays* L.) breeding lines derived from a low nitrogen maize population, 221

KAMARA, A. Y., MENKIR, A., BADU-APRAKU, B. and IBIKUNLE, O. The influence of drought stress on growth, yield and yield components of selected maize genotypes, 43

KANG'ARA, J. N. N. *see* TUWEI, P. K.

KIM, E. J., SCOLLAN, N. D., DHANOA, M. S. and BUTTERY, P. J. Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake, 103

KLING, J. G. *see* KAMARA, A. Y.

LABUSCHAGNE, M. T. *see* SOLOMON, K. F.

LOCK, A. A. The origin and significance of an indent on wheat leaves, 179

LOMBNÆS, P. and SINGH, B. R. Predicting Zn and Cu status in cereals – potential for a multiple regression model using soil parameters, 349

LOZANO, A. *see* CENTENO, C.

LUND, P. *see* SORENSEN, P.

LUTMAN, P. J. W., FREEMAN, S. E. and PEKRUN, C. The long-term persistence of seeds of oilseed rape (*Brassica napus*) in arable fields. 231

MCFARLANE, N. M., CIAVARELLA, T. A. and SMITH, K. F. The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development. 241
MCKENZIE, B. A. *see* ANWAR, M. R.
MCMASTER, G. S. and WILHELM, W. W. Phenological responses of wheat and barley to water and temperature: improving simulation models. 129
MARTIN, A. *see* BARI, A.
MATHIESEN, S. D. *see* STOREHEIER, P. V.
MENKIR, A. *see* KAMARA, A. Y.
MIRALLES, D. J., SLAFER, G. A. and RICHARDS, R. A. Influence of 'historic' photoperiod during stem elongation on the number of fertile florets in wheat. 155
MIRALLES, D. J., SLAFER, G. A., RICHARDS, R. A. and RAWSON, H. M. Quantitative developmental response to the length of exposure to long photoperiod in wheat and barley. 159
MUELLER-HARVEY, I. *see* TUWEI, P. K.

NGUGI, F. K. *see* TUWEI, P. K.

PADULOSI, S. *see* BARI, A.
PAHKALA, K. *see* HAKALA, K.
PANNANGPETCH, K. *see* BANTERNG, P.
PATANOTHAI, A. *see* BANTERNG, P.
PEKRUN, C. *see* LUTMAN, P. J. W.
POOLE, J. *see* TUWEI, P. K.
PRASAD, R. *see* SHARMA, S. N.

RAWSON, H. M. *see* MIRALLES, D. J.
REUTTER, G. *see* EIMERT, K.
RICHARDS, R. A. *see* MIRALLES, D. J.

SAIN, R. S. *see* SHARMA, S. N.
SCOLLAN, N. D. *see* KIM, E. J.
SHARMA, R. K. *see* SHARMA, S. N.
SHARMA, S. N. and PRASAD, R. Yield and P uptake by rice and wheat grown in a sequence as influenced by phosphate fertilization with diammonium phosphate and Mussoorie rock phosphate with or without crop residues and phosphate solubilizing bacteria. 359
SHARMA, S. N., SAIN, R. S. and SHARMA, R. K. The genetic control of flag leaf length in normal and late sown durum wheat. 323
SHIEL, P. J. *see* HAINES, M. M.
SHUKLA, A. K. *see* YADAV, R. L.
SINGH, B. R. *see* LOMBNÆS, P.
SINGH, V. K. *see* YADAV, R. L.
SLAFER, G. A. *see* APPENDINO, M. A.
SLAFER, G. A. *see* MIRALLES, D. J.
SMITH, K. F. *see* MCFARLANE, N. M.
SOLOMON, K. F. and LABUSCHAGNE, M. T. Variation in water use and transpiration efficiency among durum wheat genotypes grown under moisture stress and non-stress conditions. 31
SORENSEN, P., WEISBJERG, M. R. and LUND, P. Dietary effects on the composition and plant utilization of nitrogen in dairy cattle manure. 79
SPINK, J. H. *see* BERRY, P. M.
STEWART, J. L. *see* TUWEI, P. K.
STOREHEIER, P. V., VAN OORT, B. E. H., SUNDSET, M. A. and MATHIESEN, S. D. Food intake of reindeer in winter. 93
STROLKA, B. *see* EIMERT, K.
STÛTZEL, H. *see* KAGE, H.
SUNDSET, M. A. *see* STOREHEIER, P. V.
SUNG, J. M. *see* JENG, T. L.

TUWEI, P. K., KANG'ARA, J. N. N., STEWART, J. L., POOLE, J., NGUGI, F. K. and MUELLER-HARVEY, I. Factors affecting biomass production and nutritive value of *Calliandra calothyrsus* leaf as fodder for ruminants. 113

VAN OORT, B. E. H. *see* STOREHEIER, P. V.
VIVEROS, A. *see* CENTENO, C.

WANG, C. S. *see* JENG, T. L.
WEINER, J. Ecology – the science of agriculture in the 21st century. 371
WEISBJERG, M. R. *see* SORENSEN, P.
WHITE, E. M. *see* BROWNE, R. A.
WILHELM, W. W. *see* MCMASTER, G. S.

YADAV, R. L., SINGH, V. K., DWIVEDI, B. S. and SHUKLA, A. K. Wheat productivity and N use-efficiency as influenced by inclusion of cowpea as a grain legume in a rice–wheat system. 213

ZAHEDI, M. and JENNER, C. F. Analysis of effects in wheat of high temperature on grain filling attributes estimated from mathematical models of grain filling. 203

Index of Subjects

Abiotic
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETR1 (Haines *et al.*), 333

Acid phosphatase
Effect of several germination conditions on total P, phytate P, phytase, acid phosphatase activities and inositol phosphate esters in spring and winter wheat (Centeno *et al.*), 313

Agriculture
Ecology - the science of agriculture in the 21st century (Weiner), 371

Alleles
Earliness *per se* and its dependence upon temperature in diploid wheat lines differing in the major gene *Eps-A^{m}1* alleles (Appendino and Slafer), 149

Arabidopsis
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETR1 (Haines *et al.*), 333

Asparagus officinalis
Fast and reliable detection of doubled-haploids in *Asparagus officinalis* by stringent RAPD-PCR (Eimert *et al.*), 73

Bacteria
Yield and P uptake by rice and wheat grown in a sequence as influenced by phosphate fertilization with diammonium phosphate and Mussoorie rock phosphate with or without crop residues and phosphate solubilizing bacteria (Sharma and Prasad), 359

Barley
Phenological responses of wheat and barley to water and temperature: improving simulation models (McMaster and Wilhelm), 129
Quantitative developmental response to the length of exposure to long photoperiod in wheat and barley (Miralles *et al.*), 159

Biomass
Factors affecting biomass production and nutritive value of *Calliandra calothyrsus* leaf as fodder for ruminants (Tuwei *et al.*), 113
The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development (McFarlane *et al.*), 241

Biotic
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETR1 (Haines *et al.*), 333

Calliandra calothyrsus
Factors affecting biomass production and nutritive value of *Calliandra calothyrsus* leaf as fodder for ruminants (Tuwei *et al.*), 113

Cauliflower
Aspects of nitrogen use efficiency of cauliflower I. A simulation modelling based analysis of nitrogen availability under field conditions (Kage *et al.*), 1

Cereals
Predicting Zn and Cu status in cereals - potential for a multiple regression model using soil parameters (Lombnæes and Singh), 349

Concentrates
Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake (Kim *et al.*), 103

Cowpea
Wheat productivity and N use-efficiency as influenced by inclusion of cowpea as a grain legume in a rice–wheat system (Yadav *et al.*), 213

Cu
Predicting Zn and Cu status in cereals - potential for a multiple regression model using soil parameters (Lombnæes and Singh), 349

Dairy cattle
Dietary effects on the composition and plant utilization of nitrogen in dairy cattle manure (Sørensen *et al.*), 79

Development
Seasonal variation in the dynamic growth and development traits of peanut lines (Banterng *et al.*), 51
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETR1 (Haines *et al.*), 333

Diammonium phosphate
Yield and P uptake by rice and wheat grown in a sequence as influenced by phosphate fertilization with diammonium phosphate and Mussoorie rock phosphate with or without crop residues and phosphate solubilizing bacteria (Sharma and Prasad), 359

Doubled-haploids
Fast and reliable detection of doubled-haploids in *Asparagus officinalis* by stringent RAPD-PCR (Eimert *et al.*), 73

Drought stress
The influence of drought stress on growth, yield and yield components of selected maize genotypes (Kamara *et al.*), 43

Durum wheat
Variation in water use and transpiration efficiency among durum wheat genotypes grown under moisture stress

and non-stress conditions (Solomon and Labuschagne), 31
The genetic control of flag leaf length in normal and late sown durum wheat (Sharma *et al.*), 323
Fertile florets
Influence of 'historic' photoperiod during stem elongation on the number of fertile florets in wheat (Miralles *et al.*), 155
Flag leaf
The genetic control of flag leaf length in normal and late sown durum wheat (Sharma *et al.*), 323
Fractals
Use of fractals and moments to describe olive cultivars (Bari *et al.*), 63
Grain
Analysis of effects in wheat of high temperature on grain filling attributes estimated from mathematical models of grain filling (Zahedi and Jenner), 203
Effects of grain position on the panicle on starch biosynthetic enzyme activity in developing grains of rice cultivar Tainung 67 and its NaN_3-induced mutant (Jeng *et al.*), 303
Grass silage
Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake (Kim *et al.*), 103
Growth
Seasonal variation in the dynamic growth and development traits of peanut lines (Banterng *et al.*), 51
Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake (Kim *et al.*), 103
The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development (McFarlane *et al.*), 241
Phenology and growth response to irrigation and sowing date of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 273
Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 285
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETR1 (Haines *et al.*), 333
Indent
The origin and significance of an indent on wheat leaves (Lock), 179
Irrigation
Phenology and growth response to irrigation and sowing date of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 273
The effect of irrigation and sowing date on crop yield and yield components of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperature subhumid climate (Anwar *et al.*), 259
Kabuli chickpea
The effect of irrigation and sowing date on crop yield and yield components of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperature subhumid climate (Anwar *et al.*), 259
Phenology and growth response to irrigation and sowing date of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 273
Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 285
Latitudes
Comparison of central and northern European winter rye cultivars grown at high latitudes (Hakala and Pahkala), 169
Legume
Wheat productivity and N use-efficiency as influenced by inclusion of cowpea as a grain legume in a rice–wheat system (Yadav *et al.*), 213
Maize
The influence of drought stress on growth, yield and yield components of selected maize genotypes (Kamara *et al.*), 43
Agronomic performance of maize (*Zea mays* L.) breeding lines derived from a low nitrogen maize population (Kamara *et al.*), 221
Manure
Dietary effects on the composition and plant utilization of nitrogen in dairy cattle manure (Sørensen *et al.*), 79
Mathematical models
Analysis of effects in wheat of high temperature on grain filling attributes estimated from mathematical models of grain filling (Zahedi and Jenner), 203
Metabolizable energy
Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake (Kim *et al.*), 103
Moments
Use of fractals and moments to describe olive cultivars (Bari *et al.*), 63
Nitrogen
Aspects of nitrogen use efficiency of cauliflower I. A simulation modelling based analysis of nitrogen availability under field conditions (Kage *et al.*), 1
Aspects of nitrogen use efficiency of cauliflower II. Productivity and nitrogen partitioning as influenced by N supply (Kage *et al.*), 17
Dietary effects on the composition and plant utilization of nitrogen in dairy cattle manure (Sørensen *et al.*), 79
Agronomic performance of maize (*Zea mays* L.) breeding lines derived from a low nitrogen maize population (Kamara *et al.*), 221
Effect of nitrogen, seed rate and plant growth regulator (chlormequat chloride) on the grain quality of oats (*Avena sativa*) (Browne *et al.*), 249
Nutrients
Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake (Kim *et al.*), 103
Oats
Effect of nitrogen, seed rate and plant growth regulator (chlormequat chloride) on the grain quality of oats (*Avena sativa*) (Browne *et al.*), 249

Oilseed rape
The long-term persistence of seeds of oilseed rape (*Brassica napus*) in arable fields (Lutman *et al.*), 231
Olive
Use of fractals and moments to describe olive cultivars (Bari *et al.*), 63
Panicle
Effects of grain position on the panicle on starch biosynthetic enzyme activity in developing grains of rice cultivar Tainung 67 and its NaN_3-induced mutant (Jeng *et al.*), 303
Partitioning
Aspects of nitrogen use efficiency of cauliflower II. Productivity and nitrogen partitioning as influenced by N supply (Kage *et al.*), 17
Peanut
Seasonal variation in the dynamic growth and development traits of peanut lines (Banterng *et al.*), 51
Perennial ryegrass
The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development (McFarlane *et al.*), 241
Phenology
Phenology and growth response to irrigation and sowing date of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 273
Phosphate
Yield and P uptake by rice and wheat grown in a sequence as influenced by phosphate fertilization with diammonium phosphate and Mussoorie rock phosphate with or without crop residues and phosphate solubilizing bacteria (Sharma and Prasad), 359
Photoperiod
Influence of 'historic' photoperiod during stem elongation on the number of fertile florets in wheat (Miralles *et al.*), 155
Quantitative developmental response to the length of exposure to long photoperiod in wheat and barley (Miralles *et al.*), 159
Photosynthesis
The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development (McFarlane *et al.*), 241
Physiological responses
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETRI (Haines *et al.*), 333
Phytase
Effect of several germination conditions on total P, phytate P, phytase, acid phosphatase activities and inositol phosphate esters in spring and winter wheat (Centeno *et al.*), 313
Phytate P
Effect of several germination conditions on total P, phytate P, phytase, acid phosphatase activities and inositol phosphate esters in spring and winter wheat (Centeno *et al.*), 313
Plant growth regulator
Effect of nitrogen, seed rate and plant growth regulator (chlormequat chloride) on the grain quality of oats (*Avena sativa*) (Browne *et al.*), 249
Potato
Abnormalities in growth, development and physiological responses to biotic and abiotic stress in potato (*Solanum tuberosum*) transformed with Arabidopsis ETRI (Haines *et al.*), 333
Productivity
Aspects of nitrogen use efficiency of cauliflower II. Productivity and nitrogen partitioning as influenced by N supply (Kage *et al.*), 17
RAPD-PCR
Fast and reliable detection of doubled-haploids in *Asparagus officinalis* by stringent RAPD-PCR (Eimert *et al.*), 73
Reindeer
Food intake of reindeer in winter (Storeheier *et al.*), 93
Rice
Wheat productivity and N use-efficiency as influenced by inclusion of cowpea as a grain legume in a rice–wheat system (Yadav *et al.*), 213
Effects of grain position on the panicle on starch biosynthetic enzyme activity in developing grains of rice cultivar Tainung 67 and its NaN_3-induced mutant (Jeng *et al.*), 303
Yield and P uptake by rice and wheat grown in a sequence as influenced by phosphate fertilization with diammonium phosphate and Mussoorie rock phosphate with or without crop residues and phosphate solubilizing bacteria (Sharma and Prasad), 359
Root
A comparison of root and stem lodging risks among winter wheat cultivars (Berry *et al.*), 191
The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development (McFarlane *et al.*), 241
Ruminants
Factors affecting biomass production and nutritive value of *Calliandra calothyrsus* leaf as fodder for ruminants (Tuwei *et al.*), 113
Science
Ecology – the science of agriculture in the 21st century (Weiner), 371
Seasonal variation
Seasonal variation in the dynamic growth and development traits of peanut lines (Banterng *et al.*), 51
Seed rate
Effect of nitrogen, seed rate and plant growth regulator (chlormequat chloride) on the grain quality of oats (*Avena sativa*) (Browne *et al.*), 249
Simulation modelling
Aspects of nitrogen use efficiency of cauliflower I. A simulation modelling based analysis of nitrogen availability under field conditions (Kage *et al.*), 1
Phenological responses of wheat and barley to water and temperature: improving simulation models (McMaster and Wilhelm), 129
Soil
Predicting Zn and Cu status in cereals – potential for a multiple regression model using soil parameters (Lombnæes and Singh), 349
Sowing date
The effect of irrigation and sowing date on crop yield and yield components of Kabuli chickpea (*Cicer*

arietinum L.) in a cool-temperature subhumid climate (Anwar *et al.*), 259

Starch
Effects of grain position on the panicle on starch biosynthetic enzyme activity in developing grains of rice cultivar Tainung 67 and its NaN_3-induced mutant (Jeng *et al.*), 303

Steers
Effects of supplementary concentrates on growth and partitioning of nutrients between different body components in steers fed on grass silage at similar levels of metabolizable energy intake (Kim *et al.*), 103

Stem
A comparison of root and stem lodging risks among winter wheat cultivars (Berry *et al.*), 191

Subhumid
The effect of irrigation and sowing date on crop yield and yield components of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperature subhumid climate (Anwar *et al.*), 259
Phenology and growth response to irrigation and sowing date of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 273
Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 285

Temperature
Phenological responses of wheat and barley to water and temperature: improving simulation models (McMaster and Wilhelm), 129
Analysis of effects in wheat of high temperature on grain filling attributes estimated from mathematical models of grain filling (Zahedi and Jenner), 203

Total P
Effect of several germination conditions on total P, phytate P, phytase, acid phosphatase activities and inositol phosphate esters in spring and winter wheat (Centeno *et al.*), 313

Transpiration
Variation in water use and transpiration efficiency among durum wheat genotypes grown under moisture stress and non-stress conditions (Solomon and Labuschagne), 31

Water
Phenological responses of wheat and barley to water and temperature: improving simulation models (McMaster and Wilhelm), 129

Water deficits
Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 285

Waterlogging
The effects of waterlogging on growth, photosynthesis and biomass allocation in perennial ryegrass (*Lolium perenne* L.) genotypes with contrasting root development (McFarlane *et al.*), 241

Water-use efficiency
Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 285

Wheat
Phenological responses of wheat and barley to water and temperature: improving simulation models (McMaster and Wilhelm), 129
Earliness *per se* and its dependence upon temperature in diploid wheat lines differing in the major gene *Eps-A^m1* alleles (Appendino and Slafer), 149
Influence of 'historic' photoperiod during stem elongation on the number of fertile florets in wheat (Miralles *et al.*), 155
Quantitative developmental response to the length of exposure to long photoperiod in wheat and barley (Miralles *et al.*), 159
The origin and significance of an indent on wheat leaves (Lock), 179
A comparison of root and stem lodging risks among winter wheat cultivars (Berry *et al.*), 191
Analysis of effects in wheat of high temperature on grain filling attributes estimated from mathematical models of grain filling (Zahedi and Jenner), 203
Wheat productivity and N use-efficiency as influenced by inclusion of cowpea as a grain legume in a rice–wheat system (Yadav *et al.*), 213
Effect of several germination conditions on total P, phytate P, phytase, acid phosphatase activities and inositol phosphate esters in spring and winter wheat (Centeno *et al.*), 313
Yield and P uptake by rice and wheat grown in a sequence as influenced by phosphate fertilization with diammonium phosphate and Mussoorie rock phosphate with or without crop residues and phosphate solubilizing bacteria (Sharma and Prasad), 359

Winter
Food intake of reindeer in winter (Storeheier *et al.*), 93

Winter rye
Comparison of central and northern European winter rye cultivars grown at high latitudes (Hakala and Pahkala), 169

Yield
The effect of irrigation and sowing date on crop yield and yield components of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperature subhumid climate (Anwar *et al.*), 259
Water-use efficiency and the effect of water deficits on crop growth and yield of Kabuli chickpea (*Cicer arietinum* L.) in a cool-temperate subhumid climate (Anwar *et al.*), 285

Zn
Predicting Zn and Cu status in cereals – potential for a multiple regression model using soil parameters (Lombnæes and Singh), 349

21st century
Ecology – the science of agriculture in the 21st century (Weiner), 371